

10/01 9482

SEP 2000 (21 09.00)

#2
4

C400/773

PA 287378

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

August 17, 2000

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/140,988

FILING DATE: June 29, 1999

REC'D 03 OCT 2000

WIPO PCT

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)



By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS

H. L. JACKSON

Certifying Officer

1c648 U.S. PTO
06/29/99

PROVISIONAL APPLICATION COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION under 37 CFR 1.53(b) (2).

Docket Number		1952/ 47985PV		Type a plus sign (+) inside this box -	+
INVENTOR(S) / APPLICANT(S)					
LAST NAME	FIRST NAME	MIDDLE INITIAL	RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY)		
PRIVE	Gil		Toronto, Canada		
TITLE OF THE INVENTION (280 characters max)					
PEPTIDE CONJUGATES FOR THE STABILIZATION OF MEMBRANE PROTEINS AND INTERACTIONS WITH BIOLOGICAL MEMBRANES					
CORRESPONDENCE ADDRESS					
J. D. Evans 1200 G Street N.W. Suite 700 Washington, D.C. 20005					
STATE	DC	ZIP CODE	20005	COUNTRY	USA
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification	Number of Pages	19	<input type="checkbox"/> Small Entity Statement		
<input checked="" type="checkbox"/> Drawing(s)	Number of Sheets	3	<input type="checkbox"/> Other (specify)		
METHOD OF PAYMENT (check one)					
<input checked="" type="checkbox"/> A check or money order is enclosed to cover the Provisional filing fees	PROVISIONAL FILING FEE AMOUNT(S)		\$150.00		
<input type="checkbox"/> The Commissioner is hereby authorized to charge filing fees and credit Deposit Account Number: <input type="text"/>					

1c541 U.S. PTO
06/29/99

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No.

☐ Yes, the name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,

SIGNATURE Edward V. Charbon Reg No. 35,478 / for

Date 06/29/99

TYPED or PRINTED NAME J. D. Evans

REGISTRATION NO.
(if appropriate)

26,269

☐ Additional inventors are being named on separately numbered sheets attached hereto

PROVISIONAL APPLICATION FILING ONLY

Title

Peptide conjugates for the stabilization of membrane proteins and interactions with biological membranes

Field of the Invention

5 This invention generally relates to compounds that have utility as detergents. In particular, the present invention relates to a novel class of peptide-based chemical compounds that interact with proteins, lipids and other molecules. The compounds may be used for the stabilization and crystallization of proteins and membrane proteins, in particular. The compounds are also useful for modifying the properties of lipid bilayer
10 membranes, and have potential uses as cytolytic agents, as molecules that can facilitate the transport of polar molecules across biological membranes, and as emulsifiers and surfactants.

Background of the Invention

Membrane proteins are critical components of all biological membranes, and can
15 function as enzymes, receptors, channels and pumps. They are also very common in biological systems, and constitute 20-40% of all proteins found in the bacteria, archaea and eukaryotes (Wallin and von Heijne, *Protein Sci*, **7**, 1029-38 (1998), Boyd, et al., *Protein Sci*, **7**, 201-5 (1998), Gerstein, *Proteins*, **33**, 518-34 (1998), Jones, *FEBS Lett*, **423**, 281-5 (1998), Arkin, et al., *Proteins*, **28**, 465-6 (1997)). Many clinically useful drugs,
20 including the widely prescribed drugs, Prozac and Prilosec, interact with human membrane proteins. However, despite the abundance and importance of membrane proteins, this class of molecules is still only poorly understood at a structural level, mainly because of difficulties in growing crystals of membrane proteins suitable for analysis by x-ray crystallography (Garavito, et al., *J Bioenerg Biomembr*, **28**, 13-27 (1996), Ostermeier
25 and Michel, *Curr Opin Struct Biol*, **7**, 697-701 (1997), Garavito, *Curr Opin Biotechnol* **9**, 344-349 (1998)).

In order to understand the mechanism of action of a particular membrane protein, it is essential to know the three-dimensional structure of the molecule to a resolution that

reveals its atomic structure. This is typically taken to be better than 0.3 nm resolution, and nearly all of the membrane protein structures that are known to this resolution have been determined by the technique of x-ray crystallography (Branden and Tooze, *Introduction to Protein Structure*, Garland Publishing Inc., New York (1998)). If the protein in question is

5 medically important, knowledge of the 3-dimensional structure of the protein is a prerequisite for the development of new therapeutics using structure-based rational drug design methodologies (Wanke and DuBose, *Pharm Pract Manag Q*, **18**, 13-22 (1998)). The techniques used in the study of membrane protein crystals are very similar to those used for crystals of soluble proteins, and the main barrier to advancement in this field is

10 the generation of diffraction-quality crystals.

The techniques used for the crystallization of membrane proteins are generally similar to the techniques used for the crystallization of soluble proteins. Typically, a purified, concentrated solution of protein is brought to the limit of its solubility over the course of days or weeks, resulting in either the formation of a protein precipitate or of protein

15 crystals. Because precipitates are more often observed than crystals, numerous conditions are tested in these trials. The number of trials can vary in number from a few dozen to several thousand in attempts to find conditions resulting in crystal formation. The tested conditions can differ in pH, nature of added salts, concentration of the added salts, nature of the precipitant, concentration of the precipitant, temperature, and other factors

20 (A. McPherson, in "Crystallization of Biological Macromolecules", Cold Spring Harbour Press (1998)). In some instances, conditions producing suitable crystals for analysis by x-ray diffraction are not discovered even after extensive screening.

If the protein under consideration is an intrinsic membrane protein, the protein sample used in the crystallization trials is first purified and stabilized in a specific detergent in

25 order to preserve the native conformation of the protein in the absence of a lipid bilayer (H. Michel, *Trends Biochem. Sci.* **8**, 56-59 (1983), W. Kuhlbrandt, *Quart. Rev. Biophysics* **21**, 429-477 (1988)). In most instances, a number of different detergents are tested for their ability to stabilize a particular membrane protein, and for their effect in the

crystallization trials. Examples of detergents suitable for these purposes include the alkyl glycoside detergents such as octyl β -D-glucopyranoside (OG) and dodecyl β -D-maltopyranoside (DDM) (Baron and Thompson, *Biochim. Biophys. Acta* **382**, 276-285 (1975), Rosevear et al., *Biochemistry* **19**, 4108-4115 (1980)), the polyoxyethylene alkyl ether detergents such as pentaethylene glycol monooctyl ether (C8E5) and octoethylene glycol monododecyl ether (C12E8) (Garavito and Rosenbusch, *Meth. Enzymol.* **125**, 309-328 (1986), Victoria and Mahan, *Biochim Biophys Acta* **644**, 226-232 (1981)), and the detergents described in U.S. Patent No. 5,674,987, which are prepared from the reaction of a cycloalkyl aliphatic alcohol and a saccharide. Detergent-solubilized membrane proteins exist as protein-detergent complexes (PDC) in which a ring of detergent molecules covers the surface of the protein that is normally embedded in the lipid bilayer. The hydrophobic portions of the detergent amphiphiles interact with the protein surfaces normally in contact with the lipid acyl chains, and thus mimic the normal lipid environment at the surface of the membrane protein. This micelle-like ring of detergent molecules surrounding the membrane protein is very dynamic and mobile, such that the surface properties of the PDC is in general poorly suited to the formation of well-ordered crystals (Crystallization of Membrane Proteins, H. Michel ed. CRC Press, Boca Raton, FL (1991)). This is unfavorable effect is lessened in cases where the protein has large extramembranous domains, or with detergents that have small micellar volumes.

A number of techniques have been developed to address this difficulty in attempts to achieve membrane protein crystallization. For example, the formation of a complex with an antibody fragment has been used to increase the polar surface area of the *Paracoccus denitrificans* cytochrome oxidase, resulting in well-diffracting crystals (Ostermeier et al., *Nat Struct Biol*, **2**, 842-6 (1995), Ostermeier et al., *Proc Natl Acad Sci USA*, **94**, 10547-53 (1997)). Fusion proteins of the membrane protein lactose permease with soluble carrier domains have been made in attempts to achieve a similar result (Privé et al., *Acta Cryst D* **50**, 375-379 (1994), Privé and Kaback, *J Bioenerg Biomembr* **28**, 29-34 (1996)). Bacteriorhodopsin has been crystallized from cubic lipid phases (Landau and

Rosenbusch, Proc Natl Acad Sci U S A, 93, 14532-5 (1996)) in a method that does not rely on detergents at all. However, the general applicability of this method to proteins other than bacteriorhodopsin remains unproven. A strategy to reduce the volume and dynamics of the detergent surface of the PDC has been proposed by Schafmeister et al. (Science, 262, 734-8 (1993)). In this approach, amphipathic peptides have been used in the place of traditional detergents such as octyl glucoside. The peptides were designed such that the peptide would form an α -helix with one hydrophilic face and one hydrophobic face. The intention was that the hydrophobic surface of the peptide would associate with the transmembrane surface of a membrane protein. Although the peptide used in this study could maintain some membrane proteins in a solubilized state for a few days, the proteins were not sufficiently stabilized for the purposes of crystallization. Because of their limited effectiveness as detergents, these peptides have not found general utility as tools for the study of membrane proteins.

In the traditional detergents consisting of a polar head group and a linear alkyl tail, the length of the hydrocarbon moiety is an important factor in determining the ability of the detergent to preserve the native conformation of a solubilized membrane protein. Within the framework of a common head group, longer chain length detergents are generally more stabilizing towards membrane proteins, and are considered to be more "gentle". The presumed mechanism for stabilization is that the longer chains are deemed to be more effective at masking the hydrophobic transmembrane surface of the membrane protein than the short chain detergents and are thus better mimics of the native membrane environment. However, longer chain detergents occupy a larger volume of the belt region of the PDC, a feature that is expected to reduce the probability of crystallization of the complex (Michel, 73-87 in "Crystallization of Membrane Proteins", H. Michel, ed., CRC Press. Boca Raton, FL (1991)). Another factor affecting the choice of a particular detergent is the solubility of the detergent in water or buffer solutions. As the alkyl chain length increases in a series of detergents with a common head group, the overall solubility of the detergent decreases, eventually to levels making the detergent impractical for most

uses. Thus, octyl glucoside is soluble to levels greater than 20% in water, while decyl glucoside is soluble to only 0.1% in similar conditions, and dodecyl glucoside is soluble only to 0.008% (Anatrace Inc., Maumee OH 1999-2000 Catalogue). With a larger head group such as maltoside, the solubility of the long chain detergents increases, but

5 solubility is still reduced to impractical levels with hexadecyl chain lengths or longer. Thus, within a series of traditional detergents, there is conflict in the preferred length of the alkyl chain length. Long chains favor protein stability, and short chains are optimal for crystallization and detergent solubility. Since protein stability is a prime concern for crystallization trials, many membrane protein crystallization trials are carried out under
10 sub-optimal conditions.

There is a need, thus, for a non-denaturing detergent which effectively mimics the membrane's lipid bilayer, is capable of solubilizing membrane proteins in such a way that the three-dimensional conformation necessary for crystallization is retained, and has features to enhance the probability of crystallization of membrane proteins.

15 **Summary of the Invention**

Accordingly, in one aspect, the present invention provides an amphipathic peptide conjugate having detergent properties and having a hydrophobic face and a hydrophilic face, said peptide moiety of the conjugate comprising a first end and a second end, wherein said first end is covalently linked to a first aliphatic hydrocarbon moiety and said
20 second end is covalently linked to a second aliphatic hydrocarbon moiety, said aliphatic moieties linked such that they associate with the peptide moiety of the conjugate.

Preferably the peptide conjugate is a lipopeptide detergent.

Generally, a purified protein in a known detergent is subjected to a process whereby the known detergent is exchanged for the novel detergent of the present invention. The
25 protein in the novel detergent may then be subjected to conditions that promote crystallization to occur.

Brief Description of the Drawings

The present invention is described in further detail herein by reference to the following drawings in which:

Figure 1A is a schematic representation of a single lipopeptide detergent (LPD) molecule in accordance with the present invention;

Figure 1B is a schematic representation of a cylindrical assembly of several lipopeptides in which the aliphatic hydrocarbon tails are clustered in the core of the assembly;

Figure 1C is a schematic representation of a membrane protein solubilized by a traditional detergent (prior art);

Figure 1D is a schematic representation of a membrane protein solubilized by a lipopeptide detergent in accordance with the present invention;

Figure 2 is a graph showing the effectiveness of 5 lipopeptide detergents in maintaining the membrane protein bacteriorhodopsin in a soluble, stable state in the absence of a phospholipid membrane; and

Figure 3 is a scatter plot demonstrating that the lipopeptide detergents interact with phospholipid membranes, dissolving them into micelles.

Detailed Description of the Invention

Detailed Description of the Drawings

Figure 1 is a schematic representation of the lipopeptide detergents. Figure 1A shows a single LPD molecule with the α -helical peptide represented in a C α tracing with grey lines, and the aliphatic acyl chains of two fatty acids coupled to side chains at either end of the peptide shown with black lines. This representation is the presumed conformation of the monomer within the assembly shown in Figure 1B. Figure 1B shows the presumed assembly of the peptides into a cylindrical assembly. The fatty acyl chains cluster in the core of the assembly, near the central axis of the cylinder. Figure 1C shows a schematic representation of a membrane protein solubilized by a traditional detergent. This is

included to show the contrast between the present invention and the prior art. Figure 1D shows a similar protein solubilized by a lipopeptide detergent.

Figures 2 and 3 are discussed in detail later in the description.

The present invention provides novel lipopeptide detergents comprising an α -helical peptide scaffold having aliphatic hydrocarbon tails covalently linked to opposite ends of the peptide scaffold.

The peptide scaffold is not particularly limited with respect to its amino acid sequence. However, the amino acid sequence is selected so as to permit formation of the peptide scaffold into an amphipathic α -helical conformation. Generally, the peptide will comprise a mixture of hydrophobic and hydrophilic regions. Hydrophobic regions will include, but are not limited to, neutral or hydrophobic amino acids such as alanine, valine, leucine, isoleucine, methionine, phenylalanine, tryptophane or amino acids that do not occur in nature. Preferably, the hydrophobic regions are alanine-rich to favor the formation of an α -helical conformation (Chakrabarty et al., Protein Sci, 3, 843-52 (1994)). The hydrophilic regions will include, but are not limited to, amino acids which are primarily hydrophilic in nature such as glutamate, lysine, glutamine, aspartate, asparagine, histidine, serine, tyrosine, threonine or amino acids that do not occur in nature. Preferably, the hydrophilic regions promote helix formation through the formation of (i,i+4) salt bridges (Marqusee and Baldwin, Proc Natl Acad Sci U S A, 84, 8898-902 (1987)). The hydrophilic regions of the peptide align on the face of the helix that will interact with bulk aqueous phase when in a lipopeptide assembly as shown in Figure 1B. The neutral or hydrophobic face will include two residues for covalent coupling of the aliphatic moieties in the peptide conjugate. These residues will be near the termini of the peptide, at positions where they are aligned with the hydrophobic face of the peptide. The two residues can be lysine, ornithine, cysteine, glutamate or aspartate residues, but are not limited to these amino acids. Preferably, the two residues are ornithines.

The number of amino acids in the peptide scaffold is variable, and will generally be selected such that the length of the peptide scaffold when in an α -helical conformation will approximate the width of a natural membrane phospholipid bilayer, i.e. between 3.0 - 4.5 nm. Accordingly, the number of amino acids in the peptide scaffold will range from about 5 15 to 35 amino acids. Preferably, the number of amino acids in the scaffold will be about 20 - 30. More preferably, the peptide scaffold will include about 25 amino acids, or a number of amino acids which when in an α -helical conformation measure a length of about 3.7 nm.

The terminal amino acids of the peptide scaffold are also selected to promote α -helix 10 formation, and may be naturally occurring amino acids or modified forms thereof. Modifications commonly made to terminal amino acids in peptides include the addition of groups conventionally used in the art of peptide chemistry which will not adversely affect the function of the lipopeptide. For example, suitable N-terminal blocking groups can be introduced by alkylation or acylation of the N-terminus. Examples of suitable N-terminal 15 blocking groups include C₁-C₈ branched or unbranched alkyl groups, acyl groups such as formyl and acetyl groups, as well as substituted forms thereof, such as the acetamidomethyl (Acm) group. Desamino analogs of amino acids are also useful N-terminal blocking groups, and can either be coupled to the N-terminus of the peptide or used in place of the N-terminal residue. Suitable C-terminal blocking groups, in which the carboxyl group of the C-terminus 20 is either incorporated or not, include esters, ketones or amides. Ester or ketone-forming alkyl groups, particularly lower alkyl groups such as methyl, ethyl and propyl, and amide-forming amino groups such as primary amines (-NH₂), and mono- and di-alkylamino groups such as methylamino, ethylamino, dimethylamino, diethylamino, methylethylamino and the like are examples of C-terminal blocking groups. Descarboxylated amino acid analogues 25 such as agmatine are also useful C-terminal blocking groups and can be either coupled to the peptide's C-terminal residue or used in place of it. Further, it will be appreciated that the free amino and carboxyl groups at the termini can be removed altogether from the peptide to yield desamino and descarboxylated forms thereof without affect on peptide function.

Preferred examples of such modifications include N-terminal acetylation and C-terminal amidation which are known to promote α -helix formation (Doig et al., Biochemistry, 33, 3396-403 (1994)).

Internal amino acids of the peptide may also be modified by derivatization provided that
5 this modification does not affect the function of the lipopeptide, and does not interfere with its α -helical conformation. Such derivatizations can be made to the side chains of the amino acids. For example, the side chains can be derivatized by incorporation of blocking groups as described above.

The peptide conjugate may be readily prepared by standard, well-established solid-
10 phase peptide synthesis (SPPS) as described by Stewart et al. in Solid Phase Peptide Synthesis, 2nd Edition, 1984, Pierce Chemical Company, Rockford, Illinois; and as described by Bodanszky and Bodanszky in The Practice of Peptide Synthesis, 1984, Springer-Verlag, New York. Other synthetic protocols, including biological or solution phase methods, can also be used. For the SPPS method, a suitably protected amino acid residue
15 is first attached through its carboxyl group to a derivatized, insoluble polymeric support, such as cross-linked polystyrene or polyamide resin. "Suitably protected" refers to the presence of protecting groups on both the α -amino group of the amino acid, and on any side chain functional groups. Side chain protecting groups are generally stable to the solvents, reagents and reaction conditions used throughout the synthesis, and are removable under
20 conditions which will not affect the final peptide product. Stepwise synthesis of the oligopeptide is carried out by the removal of the N-protecting group from the initial amino acid, and coupling thereto of the carboxyl end of the next amino acid in the sequence of the desired peptide. This amino acid is also suitably protected. The carboxyl of the incoming amino acid can be activated to react with the N-terminus of the support-bound amino acid by
25 formation into a reactive group such as formation into a carbodiimide, a symmetric acid anhydride or an "active ester" group such as hydroxybenzotriazole or pentafluorophenyl esters.

Examples of solid phase peptide synthesis methods include the BOC method which

utilizes tert-butoxycarbonyl as the α -amino protecting group, and the Fmoc method which utilizes 9-fluorenylmethoxycarbonyl to protect the α -amino of the amino acid residues, both methods of which are well-known by those of skill in the art.

The aliphatic moieties can be coupled to the resin-coupled peptide by selectively deblocking amino acid side chain protecting groups, followed by reaction with an appropriate aliphatic derivative. Aliphatic derivatives suitable for this purpose include, but are not limited to, saturated fatty acids, unsaturated fatty acids, branched fatty acids, cyclic alkyl acids, aromatic alkyl acids, alkyl amines, alkyl maleimides, alkyl acid chlorides, and alkyl anhydrides. Several strategies can be used to couple the aliphatic derivative to the peptide.

For example, if the peptide is synthesized with the Fmoc method, a BOC group can be used as the protecting group on the δ -amino group of the ornithine monomers identified as sites for aliphatic coupling. Upon completion of the synthesis of the main peptide chain, the ornithine BOC groups can be selectively removed with trifluoroacetic acid, generating free primary amino functionalities at these positions. Reaction with an aliphatic derivative such as a fatty acid can be used to form a peptide linkage with each of the two ornithine side chains. Examples of suitable saturated fatty acids include octanoic acid, nonanoic acid, decanoic acid, undecanoic acid, dodecanoic acid, tridecanoic acid, tetradecanoic acid, pentadecanoic acid, hexadecanoic acid, heptadecanoic acid, octadecanoic acid, nonadecanoic acid, eicosanoic acid, heneicosanoic acid, docosanoic acid, tricosanoic acid, tetracosanoic acid, pentacosanoic acid, and hexacosanoic acid. Following the coupling of the aliphatic groups, the remaining amino acid side chains can be deblocked under appropriate conditions, such as with hydrofluoric acid or trifluoromethanesulfonic acid.

Incorporation of N- and/or C- blocking groups can also be achieved using protocols conventional to solid phase peptide synthesis methods. For incorporation of C-terminal blocking groups, for example, synthesis of the desired peptide is typically performed using, as solid phase, a supporting resin that has been chemically modified so that cleavage from the resin results in a peptide having the desired C-terminal blocking group. To provide

peptides in which the C-terminus bears a primary amino blocking group, for instance, synthesis is performed using a p-methylbenzhydrylamine (MBHA) resin so that, when peptide synthesis is completed, treatment with hydrofluoric acid or trifluoromethanesulfonic acid (TFMSA) releases the desired C-terminally amidated peptide. Similarly, incorporation of an N-methylamine blocking group at the C-terminus is achieved using N-methylaminoethyl-derivatized DVB resin, which upon HF treatment releases a peptide bearing an N-methylamidated C-terminus. Blockage of the C-terminus by esterification can also be achieved using conventional procedures. This entails use of resin/blocking group combination that permits release of side-chain protected peptide from the resin, to allow for subsequent reaction with the desired alcohol, to form the ester function. Fmoc protecting groups, in combination with DVB resin derivatized with methoxyalkoxybenzylalcohol or equivalent linker, can be used for this purpose, with cleavage from the support being effected by TFA in dichloromethane. Esterification of the suitably activated carboxyl function e.g. with DCC, can then proceed by addition of the desired alcohol, followed by deprotection and isolation of the esterified peptide product.

Incorporation of N-terminal blocking groups can be achieved while the synthesized peptide is still attached to the resin, for instance by treatment with a suitable anhydride and nitrile. To incorporate an acetyl blocking group at the N-terminus, for instance, the resin-coupled peptide can be treated with 20% acetic anhydride in acetonitrile. The N-blocked peptide product can then be cleaved from the resin, deprotected and subsequently isolated.

To ensure that the peptide obtained from either chemical or biological synthetic techniques is the desired peptide, analysis of the peptide composition should be conducted. Such amino acid composition analysis may be conducted using high resolution mass spectrometry (MS) to determine the molecular weight of the peptide. Alternatively, or additionally, the amino acid content of the peptide can be confirmed by hydrolyzing the peptide in aqueous acid, and separating, identifying and quantifying the components of the mixture using HPLC, or an amino acid analyzer. Protein sequenators, which sequentially degrade the peptide and identify the amino acids in order, may also be used to determine

definitely the sequence of the peptide.

Having obtained the desired peptide conjugate, purification to remove contaminants is generally then conducted. Any one of a number of conventional purification procedures may be used to attain the required level of purity including, for example, ion-exchange and gel
5 filtration chromatography or reversed-phase high-pressure liquid chromatography (HPLC) using an alkylated silica column such as C₄-, C₈- or C₁₈- silica. A gradient mobile phase of increasing organic content is generally used to achieve purification, for example, acetonitrile in an aqueous buffer, usually containing a small amount of trifluoroacetic acid. Because the overall hydrophobicity of the peptide conjugates increases with larger aliphatic moieties, C₄-
10 silica is the preferred chromatographic resin for these compounds.

Aliphatic hydrocarbon moieties are linked in a covalent manner to both the N- and C-termini of the scaffold peptide or to sites near each of these termini such that they associate with the hydrophobic region of the peptide scaffold. In one embodiment of the present invention, the aliphatic hydrocarbon tails are linked to ornithine residues located
15 adjacent to N- and C- terminal alanine residues of the scaffold peptide. The δ -amino groups of the ornithines are coupled to the carboxyl groups of tetracosanoic acid via peptide linkages. Ornithines are used in place of the more common lysine residues as sites for the hydrocarbon tail linkage since they have fewer methylene groups between the main chain peptide atoms and the side chain amine, and may position the hydrocarbon
20 chains more precisely in association with the hydrophobic region of the peptide.

The lipopeptide detergent is advantageous over "traditional" detergents due to its ability to self-associate into a cylinder of defined dimensions. The cylinders are made up of colinear α -helices and themselves associate into a cylindrical assembly, as shown in Fig. 1B, in which the hydrophilic surfaces of the individual helices are exposed to the bulk
25 aqueous phase and the hydrocarbon tails are packed in the core of the assembly effectively mimicking the chains in a membrane phospholipid bilayer. Fig. 1D illustrates how a membrane protein can be accommodated in the core of a lipopeptide assembly with the aliphatic hydrocarbon tails forming a cylindrical layer against the protein, again

better mimicking biological membrane conformation, allowing for preservation of the biological activity of solubilized membrane proteins.

The described lipopeptide detergents with two coupled aliphatic moieties ranging from ten to twentyfour carbon alkyl chains are soluble in water, in contrast to an alkyl chain length maximum of sixteen carbon groups in the traditional detergents. The favorable solubility properties of the long chain lipopeptide detergents make it possible for these detergents to stabilize large hydrophobic surfaces of membrane proteins.

In addition to their stabilizing properties, the present lipopeptide detergents have been designed to favor the crystallization of membrane proteins. They lie close to the surface of the membrane protein, and are thus less obtrusive to the formation of a crystal lattice. Also, they present a rigid outer surface of α -helices. These are features that favor membrane protein crystallization (Schafmeister et al., Science, 262. 734-8 (1993) Michel, Crystallization of membrane proteins, 73-87 (1991)).

The lipopeptide detergents of the present invention may be used to crystallize membrane proteins. Generally, the method comprises solubilizing the membrane protein with a detergent, and then exposing the solubilized membrane to conditions which promote crystallization to occur.

The lipopeptide detergents are also membrane-active compounds, and can insert into phospholipid bilayers. At sufficiently high concentrations, they can disrupt the bilayers and form mixed lipid/lipopeptide micelles.

The lipopeptide detergents of the present invention have the activities of traditional detergents and hence they may be used to modulate and disrupt biological membranes, and therefore to transport polar molecules across membranes, including ions. As surface active agents or emulsifiers, they may be used in protein and/or lipid emulsions. They may also be used as cytolytic agents.

EXAMPLES

Embodiments of the present invention are described in further detail by reference to the following specific examples, which are not to be construed as limiting the appended claims.

5 Example 1: Synthesis of LPD-24

The lipopeptide, LPD-24, exemplifies a lipopeptide detergent in accordance with the present invention. The scaffold peptide of LPD-24 has the following chemical structure:



wherein A is alanine, O is ornithine, E is glutamate, K is lysine, and Y is tyrosine,

- 10 CH₃CONH- is the acetylated amino terminal group of the peptide, and -CONH₂ is the carboxamide end of the peptide chain. A single tyrosine is included to allow spectrophotometric detection of the peptide at 280 nm.

- LPD-24 is synthesized on a solid support resin using a combination of tert-butoxycarbonyl (Boc) and 9-Fluorenylmethoxycarbonyl (Fmoc) chemistries (Novabiochem
- 15 97/98 Catalogue and Peptide synthesis handbook). The synthesis proceeds from the C-terminus of the peptide to the N-terminus, with all the main chain peptide synthesis couplings based on Fmoc chemistry. The resin tert-butoxycarbonyl-Alanine-methylbenzhydrylamine (Boc-Ala-MBHA) is chosen so as to produce a peptide carboxamide upon cleavage from the resin. The resin is first prepared by removal of the
- 20 Boc protecting group with 50% trifluoroacetic acid (TFA), generating the free α -amino acid amine of the alanine. Sequential addition of the following 24 amino acids proceeds with the coupling of the appropriate Fmoc-protected amino acid: N- α -Fmoc-L-alanine (Fmoc-Ala), N- α -Fmoc-L-glutamamic acid α -benzyl ester (Fmoc-Glu(Obz)), N- α -Fmoc-N- α -2-chloro-benzyloxycarbonyl-L-lysine (Fmoc-Lys(2ClZ)), N- α -Fmoc-N- α -tert-butoxycarbonyl-
- 25 L-ornithine (Fmoc-Om(Boc)), or N- α -Fmoc-O-2-bromo-benzyloxycarbonyl-L-tyrosine (Fmoc-Tyr(2BrZ)) with the coupling reagent O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluroniumhexafluorophosphate (HATU). Upon completion of the coupling

reaction, the Fmoc protecting group is removed with 20% piperidine in preparation for the next amino acid coupling. Following the addition of the last amino acid and the removal of the Fmoc group, the amine terminus of the chain is acetylated with acetic anhydride.

Next, the Boc protecting groups of the ornithine side chains are removed with 50% TFA in preparation for the coupling with the fatty acid. Two equivalents of tetracosanoic acid are coupled to the peptide with HATU. The final step involves the cleavage of the peptide from the resin and the deprotection of the glutamate, lysine, and tyrosine side chains with trifluoromethanesulfonic acid (TFMSA).

The lipopeptide is precipitated in ether, and washed four times in ether. The white pellet is dissolved in water, lyophilized, and redissolved in water. The peptide is purified by gel filtration chromatography in ammonium carbonate buffer, lyophilized, and redissolved in water. The correctness of the synthesis is confirmed by electrospray ionization mass spectrometry (ESI-MS).

Lipopeptide detergents with pairs of aliphatic hydrocarbon tails of length 10, 12, 14, 16, 18, 20, 22, 24, and 28 carbons (LPD-10 to LPD-28) have been designed and synthesized. The LPDs with chain lengths up to 24 carbons are soluble in water to over 50 mg/ml. LPD-28 is poorly soluble in water. Computer-assisted molecular modelling suggests that the longer chains (C16 and greater) can cross past each other and that each may span the entire length of the helix. Every batch of peptide is analyzed by ESI-MS to confirm the synthesis. The uncoupled peptide is referred to as LPD-0, and does not have detergent properties.

Example 2: Effectiveness of Lipopeptides in stabilizing solubilized membrane proteins

Bacteriorhodopsin was purified from purple membranes (isolated from *Halobacterium halobium*, a generous gift from J. Lanyi, UC Riverside). 4 mL of purple membranes were diluted with 10 ml 1% Triton X100, 10 mM NaH₂PO₄ pH 5.7, and mixed for 12 hours at 30°C in the dark. The solution was then centrifuged for 15 minutes at 55,000 r.p.m in a Beckman NVT 90 centrifuge rotor to remove any insoluble material. The supernatant was

added to a diethylaminoethyl (DEAE) sepharose ion exchange column (10 ml column volume) previous equilibrated with 1% Triton X100, 10 mM NaH₂PO₄ pH 5.7. The column was then washed with 25 mL 34 mM OG, 10 mM NaH₂PO₄ pH 5.7, followed by elution with a buffer containing 34 mM OG, 500 mM NaH₂PO₄ pH 5.7. OG is a standard
5 detergent for the purification and crystallization of bacteriorhodopsin (G.F. Schertler et al., J. Mol. Biol **234**, 156-164 (1993)). LPD (lipopeptide) detergents (LPD-12, LPD-14, LPD-16, LPD-18 and LPD-20) were added to the protein solutions at a final concentration of 3.25 mM, and the OG was removed by dialysis in the dark through membranes with molecular weight cutoffs of 1 kDa. The LPD detergents were fully retained in the protein
10 solution, while the OG was removed from the protein solution to levels of less than 0.5 mM after 2 days. The samples were stored in the dark at room temperature, and at 1 day, 4 days, 12 days, 18 days and 25 days storage, the samples were centrifuged at 100,000 X g for 45 minutes and the absorbance at 540 nm of the supernatants were measured. Solubilized, properly folded, native bacteriorhodopsin remains in the supernatant and has
15 an absorbance maximum at 540 nm. All LPDs with chain lengths of 12 amino acids and longer were much more stabilizing to the membrane protein bacteriorhodopsin than octyl glucoside (OG) as illustrated in Fig. 2.

Example 3: Interaction of lipopeptide detergents with phospholipid membranes

20 Phosphatidyl choline vesicles were prepared by extrusion through 0.1 nm pore membranes (Avestin, Ottawa), at 1 mM concentration, and diluted to 0.1 mM phospholipid. Dodecyl maltoside (DDM) or lipopeptide detergent were added to the indicated concentrations, and the hydrodynamic radius of the suspension was measured on a DynaPro-800 dynamic light scattering device.

25 The results of this experiment are illustrated in Fig. 3. This scatter plot demonstrates that the lipopeptide detergents interact with phospholipid membrane vesicles, dissolving them into micelles. The control C-0 peptide had no effect on these vesicles.

The disclosure of all of the literature references and any patents referred to herein are incorporated herein by reference.

While the invention has been described with particular reference to certain embodiments thereof, it will be understood that those of ordinary skill in the art within the scope and spirit of the following claims may make changes and modifications.

In the claims, the word "comprising" means "including the following elements (in the body), but not excluding others"; the phrase "consisting of" means "excluding more than traces of other than the recited ingredients"; and the phrase "consisting essentially of" means "excluding unspecified ingredients which materially affect the basic characteristics of the composition".

I Claim:

1. An amphipathic peptide conjugate having detergent properties and having a hydrophobic face and a hydrophilic face, said peptide moiety of the conjugate comprising a first end and a second end, wherein said first end is covalently linked to a first aliphatic hydrocarbon moiety and said second end is covalently linked to a second aliphatic hydrocarbon moiety, said aliphatic moieties linked such that they associate with the peptide moiety of the conjugate.
2. The peptide conjugate as defined in claim 1, which comprises a lipopeptide detergent.
3. The peptide conjugate as defined in claim 1, wherein said peptide comprises hydrophobic and hydrophilic regions.
4. The peptide conjugate as defined in claim 1, wherein said peptide comprises 15-35 amino acids.
5. The peptide conjugate as defined in claim 4, wherein said peptide comprises about 25 amino acids.
6. The peptide conjugate as defined in claim 5, wherein said peptide has the amino acid sequence, AOAEAAEKA AKYAAEAAEKA AKAOA.
7. The peptide conjugate as defined in claim 1, wherein the length of said peptide is approximately equal to the width of a phospholipid bilayer.
8. The peptide conjugate as defined in claim 7, wherein the length of said peptide is in the range of about 3.5 - 4.0 nm.
9. The peptide conjugate as defined in claim 8, wherein the length of said peptide is about 3.7 nm.
10. The peptide conjugate as defined in claim 1, wherein the termini of said peptide are protected.
11. The peptide conjugate as defined in claim 10, wherein the N-terminus of said peptide is acetylated and the C-terminus of said peptide is amidated.
12. The peptide conjugate as defined in claim 1, wherein said aliphatic hydrocarbon moieties comprise from about 8-24 carbon atoms.

ABSTRACT

The present invention provides a novel class of detergents referred to herein as lipopeptide detergents. Lipopeptide detergents comprise an amphipathic α -helical peptide having a hydrophobic or neutral face and a hydrophilic face. To both ends of this peptide
5 is covalently linked aliphatic hydrocarbon tails, these aliphatic tails being linked therein such that they associate with the hydrophobic or neutral face of the peptide. Lipopeptide detergents can advantageously be used to solubilize membrane proteins in a manner that permits the subsequent crystallization thereof.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215




Figure 1




Figure 2.




Figure 3.